

Microwave Spectrum and Ground State Energy Levels of H_2^{17}O

FRANK C. DE LUCIA AND PAUL HELMINGER¹

Department of Physics, Duke University, Durham, North Carolina 27706

The rotational spectrum of H_2^{17}O in the ground vibrational state has been investigated by means of high resolution microwave spectroscopy. Fifteen transitions in the 2.0-0.4 mm wavelength region have been measured. Both the quadrupole and spin-rotation molecular hyperfine tensors for the ^{17}O nucleus are calculated from the observed hyperfine splittings of the rotational transitions. In addition, several of the low- J lines are of significance in the atmospheric absorption of electromagnetic radiation in the microwave region. The observed microwave transitions are combined with our earlier analyses of H_2^{16}O and H_2^{18}O to calculate the ground state energy levels of H_2^{17}O .

I. INTRODUCTION

Although isotopic species of water have been extensively studied for many years in the infrared and more recently in the microwave region (1), little work has been reported on H_2^{17}O . Infrared studies have been difficult because of the overlapping bands of H_2^{16}O and H_2^{18}O which result from the high concentration of these species in isotopically enriched samples of H_2^{17}O . The microwave spectrum of H_2^{17}O consists of only two rotational transitions at wavelengths longer than 1 mm, at 13 GHz and at 194 GHz (2). We have observed the latter under high resolution and with good signal-to-noise and have resolved seven hyperfine components which result from the quadrupole moment of the ^{17}O nucleus. In addition, we have observed the hyperfine components of five submillimeter transitions. These measurements are now possible because of both a significant increase in the spectral region accessible to our microwave spectrometer and a corresponding order of magnitude increase in sensitivity in the region of the submillimeter spectrum previously available to us (3). Among the newly observed lines are several low- J transitions which make significant contributions to the atmospheric absorption in the submillimeter region even though the natural abundance of ^{17}O is only 0.04%. These measurements, when combined with the results of our previous analyses of H_2^{16}O (4) and H_2^{18}O (5), make possible the analysis of H_2^{17}O without the explicit inclusion of H_2^{17}O infrared data. The energy levels which result from this analysis are in good agreement with combination differences which can be calculated from lines of H_2^{17}O observed by Fraley, Rao, and Jones (6) and by Williamson and Rao (7) as impurities in infrared H_2^{18}O spectra.

¹ Present address: Department of Physics, University of South Alabama, Mobile, Alabama 36688. This work was supported by NSF Grant GP-34590.

Table I. Hyperfine Interaction Constants for ^{17}O in H_2^{17}O .

<u>Spectral Constants</u>		
	<u>3_{13}</u>	<u>2_{20}</u>
eQq_J	$4.258 \pm .020 \text{ MHz}$	$-3.000 \pm .020 \text{ MHz}$
C	$-0.012 \pm .003 \text{ MHz}$	$-0.017 \pm .003 \text{ MHz}$
<u>Molecular Constants</u>		
<u>Quadrupole</u>	<u>Spin-Rotation ($\times 10^9$)</u>	
$V_{aa} = -8.899 \text{ MHz}$	$A_{aa} = 63.4$	
$V_{bb} = -1.269 \text{ MHz}$	$A_{bb} = 29.1$	
$V_{cc} = 10.169 \text{ MHz}$	$A_{cc} = 30.6$	

II. EXPERIMENTAL DETAILS

The millimeter and submillimeter spectrometer used for this work has been described previously (8). Briefly, a crystal harmonic generator is driven by a phase-locked OKI 55V11 klystron to produce tunable microwave power in the 100–800 GHz region. This power is directed with tapered horn and teflon lens optics through a 1-m quasifree space absorption cell and detected by a 1.5K InSb photodetector. All frequency measurements are referenced to WWVB. The absorption cell was typically filled to 50 μm with a sample isotopically enriched to 30% H_2^{17}O . This sample was kindly supplied by Drs. H. W. Morgan and P. A. Staats of Oak Ridge National Laboratory.

Fraley, Rao, and Jones (6) have proposed the empirical relation

$$[\nu(\text{H}_2^{16}\text{O}) - \nu(\text{H}_2^{17}\text{O})]/[\nu(\text{H}_2^{16}\text{O}) - \nu(\text{H}_2^{18}\text{O})] = 0.529. \quad (1)$$

This relation and the results of our previous work on H_2^{16}O and H_2^{18}O were used to initially locate the 1_{10} – 1_{01} and 2_{11} – 2_{02} transitions of H_2^{17}O at 552 and 748 GHz, respectively. These transitions, the two previously observed microwave transitions, and distortion constants calculated from the other species made possible the calculation of an initial set of rotational constants α , β , \mathcal{C} , as well as the accurate prediction of the remaining microwave lines.

III. HYPERFINE ANALYSIS

Each of the rotational transitions of H_2^{17}O is split into a number of components because of the interaction of the quadrupole moment of the ^{17}O nucleus and the gradient of the molecular electric field. Under the assumption that the difference between the equilibrium hyperfine constants and the hyperfine constants of the ground vibrational state is small, the quadrupole eQq_J and spin-rotation C constants are related to the fundamental molecular tensors \mathbf{V} and \mathbf{A} by

$$q_J = \frac{2}{(J+1)(2J+3)} \sum_a V_{aa} \langle P_a^2 \rangle, \quad (2)$$




FIG. 1. Observed hyperfine structure of the $3_{13}-2_{20}$ transition of H_2^{17}O .

$$C = g_I \frac{1}{J(J+1)} \sum_g \Lambda_{gg} B_g \langle P_g^2 \rangle, \quad (3)$$

where g refers to the principal axes of the moment of inertia tensor, P_g is the component




FIG. 2. Theoretical hyperfine structure of the $3_{13}-2_{20}$ transition of H_2^{17}O . Units are in MHz.

Table II. Observed Microwave Spectrum of H_2^{17}O (MHz).

Transition (F' ← F)	Observed	Calculated	Obs- Cal
$1_{10} \leftarrow 1_{01}$	552020.960	552020.765	0.195
$2_{11} \leftarrow 2_{02}$	748458.254	748458.368	-0.114
$3_{13} \leftarrow 2_{20}$			
$9/2 \leftarrow 7/2$	193999.796	193999.790	0.006
$7/2 \leftarrow 5/2$	194000.310	194000.274	0.036
$5/2 \leftarrow 3/2$	194001.675	194001.656	0.019
$3/2 \leftarrow 1/2$	194002.998	194002.981	0.017
$11/2 \leftarrow 9/2$	194004.172	194004.151	0.021
$9/2 \leftarrow 7/2$	194005.835	194005.815	0.020
$7/2 \leftarrow 5/2$	194007.002	194006.987	0.015
$4_{14} \leftarrow 3_{21}$			
lower	385784.860	385785.234	-0.374
upper	385788.189	385788.326	-0.137
$4_{23} \leftarrow 3_{30}$			
lower	469808.134	469808.170	-0.036
upper	469811.064	469811.045	0.019
$5_{15} \leftarrow 4_{22}$			
lower	323825.089	323824.944	0.145
upper	323827.873	323827.748	0.125
$6_{16} \leftarrow 5_{23}^a$			
$13/2 \leftarrow 11/2$	13533.79	13533.83	-0.04
$15/2 \leftarrow 13/2$	13534.41	13534.50	-0.09
$11/2 \leftarrow 9/2$	13536.02	13536.10	-0.08
$9/2 \leftarrow 7/2$	13537.06	13537.09	-0.03
$17/2 \leftarrow 15/2$	13537.97	13538.00	-0.03

^aRef. 2.

of the rotational angular momentum along the g axis, B_g is the rotational constant for the g -axis, and g_I is the nuclear g factor. The products $g_I \Lambda_{gg} B_g$ are the diagonal elements of the spin-rotation tensor C_{gg} .

A precise molecular beam maser measurement of the hyperfine structure of the 2_{20} - 2_{21} transition of HD^{17}O has resulted in accurate values of $(eQq_I)^{17}\text{O}$ and $(C)^{17}\text{O}$ for these states (9). Because of the molecular symmetry of the ^{17}O site, only the diagonal elements of \mathbf{V} and $\mathbf{\Lambda}$ are nonzero. Although the Λ_{ii} cannot be calculated from these data, the V_{ii} can be by the inclusion of the Laplace condition $\Sigma V_{ii} = 0$.

Since the contribution of the spin-rotation interaction is small for most of the transitions reported here, Eq. (2) and the V_{ii} of (9) make possible a good approximation to most of the hyperfine structure observed in this work. However, an analysis of the extensive hyperfine structure of the 3_{13} - 2_{20} transition requires a consideration of spin-rotation effects. The hyperfine constants which result from this analysis are shown in Table I and the spectrum calculated from these constants is compared to the experimental spectrum in Figs. 1 and 2. These hyperfine constants can be combined with those of (9) to calculate both the V_{ii} and Λ_{ii} which are also shown in Table I. These elements of the molecular tensors will reproduce the hyperfine constants of both experiments to within their quoted error limits.

Table III. Rotational Constants of H_2^{17}O (MHz).

$A = 830282.838 \pm 8.7^{\text{a,b}}$ $B = 435341.662 \pm 8.0$ $C = 277510.396 \pm 8.0$		
$\Delta_J = 37.44401^{\text{c}}$ $\delta_J = 15.22675$	$\Delta_{JK} = -171.5584$ $\delta_K = 36.2566$	$\Delta_K = 965.7665 \pm 2.3$
$H_J = 1.48878 \times 10^{-2}$ $H_K = 4.409069 \times 10^0$	$H_{JK} = -2.10405 \times 10^{-2}$ $h_J = 7.844745 \times 10^{-3}$ $h_K = 5.4855 \times 10^{-1}$	$H_{KJ} = -4.81054 \times 10^{-1}$ $h_{JK} = -2.11625 \times 10^{-2}$
$L_{JK} = 1.53235 \times 10^{-3}$	$L_{KKJ} = 5.1476 \times 10^{-3}$ $\ell_K = 4.982 \times 10^{-3}$	$L_K = -5.966754 \times 10^{-2}$
$P_K = 7.50542 \times 10^{-4}$	$p_K = 1.88015 \times 10^{-5}$	

- a. Errors quoted are twice the statistical standard deviation and reflect uncertainties in and correlations among the distortion constants calculated from H_2^{16}O and H_2^{18}O data.
- b. The number of places quoted are required to reproduce the observed microwave lines. Because of the correlation among the distortion constants this number can be substantially larger than the uncertainty for many of the higher order constants.
- c. All distortion constants except Δ_K are calculated from H_2^{16}O and H_2^{18}O data.

IV. ROTATIONAL ANALYSIS

It has been shown that it is possible to characterize the spectra of light asymmetric rotors to within the expected experimental uncertainty of microwave measurements (~ 0.1 MHz) (10, 11) by use of Watson's formulation (12, 13, 14) of a reduced centrifugal distortion Hamiltonian. For H_2^{16}O this Hamiltonian has the form

$$\mathcal{H} = \mathcal{H}_r + \mathcal{H}_d^{(4)} + \mathcal{H}_d^{(6)} + \mathcal{H}_d^{(8)} + \mathcal{H}_d^{(10)}, \quad (4)$$

$$\mathcal{H}_r = \frac{1}{2}(\mathcal{B} + \mathcal{C})P^2 + [\mathcal{A} - \frac{1}{2}(\mathcal{B} + \mathcal{C})](P_z^2 - b_p P_-^2), \quad (4a)$$

$$\mathcal{H}_d^{(4)} = -\Delta_J P^4 - \Delta_{JK} P^2 P_z^2 - \Delta_K P_z^4 - 2\delta_J P^2 P_-^2 - \delta_K (P_z^2 P_-^2 + P_-^2 P_z^2), \quad (4b)$$

$$\begin{aligned} \mathcal{H}_d^{(6)} = & H_J P^6 + H_{JK} P^4 P_z^2 + H_{KJ} P^2 P_z^4 + H_K P_z^6 + 2h_J P^4 P_-^2 \\ & + h_{JK} P^2 (P_z^2 P_-^2 + P_-^2 P_z^2) + h_K (P_z^4 P_-^2 + P_-^2 P_z^4), \end{aligned} \quad (4c)$$

$$\mathcal{H}_d^{(8)} = L_{JK} P^4 P_z^4 + L_{KKJ} P^2 P_z^6 + L_K P_z^8 + l_K (P_z^6 P_-^2 + P_-^2 P_z^6), \quad (4d)$$

$$\mathcal{H}_d^{(10)} = P_K P_z^{10} + p_K (P_z^8 P_-^2 + P_-^2 P_z^8), \quad (4e)$$

where $P^2 = (P_x^2 + P_y^2 + P_z^2)$, $P_-^2 = (P_x^2 - P_y^2)$, and $b_p = (\mathcal{C} - \mathcal{B})/(2\mathcal{A} - \mathcal{B} - \mathcal{C})$ is Wang's asymmetry parameter appropriate for a prolate top. A similar Hamiltonian is appropriate for H_2^{18}O . For both of these species, the number of adjustable parameters exceeds the number of observed microwave transitions. As a result, the rather extensive body of infrared data for these species was used in a weighted least-squares analysis with the microwave data (4, 5).

The infrared work on H_2^{17}O is significantly less extensive and other techniques are required. Although the observed microwave transition frequencies of particular transitions of H_2^{16}O and H_2^{18}O differ by as much as 100 GHz, the distortion constants of the two species are very similar. In many cases the difference between the constants is sufficiently small that their uncertainties overlap. Since the different data sets which were used in the two analyses resulted in the inclusion of somewhat different Hamiltonian terms and since these terms are highly correlated, the similarity between the distortion in the two species is probably even greater than indicated by this comparison. Therefore, all or most of the distortion parameters can be fixed at the average of their values for the H_2^{16}O and H_2^{18}O species.

Table II shows the measured frequencies of the H_2^{17}O microwave transitions. Extensive hyperfine structure was resolved for the $3_{13}-2_{20}$ and $6_{16}-5_{23}$ transitions, three of the other R -branch transitions were observed as doublets, and the Q -branch transitions were observed as unresolved singlets. Unsplit rotational frequencies were calculated for the multiplet transitions by the subtraction of the hyperfine contributions from the observed frequencies. It was found that these frequencies could be fit to within the expected uncertainties of the microwave measurements by use of the Hamiltonian of Eq. (4) with Δ_K and \mathcal{A} , \mathcal{B} , \mathcal{C} as adjustable parameters. The remaining parameters were fixed at the values calculated from our earlier analyses of H_2^{16}O and H_2^{18}O . These parameters are shown in Table III. Since the values of these constants depend upon the distortion constants of H_2^{16}O and H_2^{18}O , the uncertainty in these constants can be no smaller than the uncertainty in the corresponding constants of H_2^{16}O and H_2^{18}O . Since the rotation-distortion parameters of H_2^{18}O are somewhat less certain than those of H_2^{16}O , the uncertainties in the H_2^{17}O parameters are taken to be those of H_2^{18}O .

The rotational constants \mathcal{A} , \mathcal{B} , \mathcal{C} of the Watson formulation contain contributions from the distortion term R_6 and thereby differ from the A , B , C of Kivelson and Wilson (15). Similarly, the A , B , C of Kivelson and Wilson contain contributions from other distortion terms. The removal of these contributions results in A' , B' , C' which are simply related to the moments of inertia by $I_a = h/8\pi^2 A'$, etc. The relations among these constants are discussed in (16) and Table IV shows their values for H_2^{17}O .

V. ENERGY LEVELS

Because of the large number of resolvable lines in the infrared spectra of light asymmetric molecules, their energy levels are usually constructed by a bootstrap technique which builds upon the ground state by use of measured energy differences. This approach is not possible for H_2^{17}O because of the small number of infrared measurements reported

Table IV. Rotational Constants of H_2^{17}O (MHz).

A: B: C:	830282.8	435341.7	277510.4
A: B: C:	830338.4	435146.5	277650.1
A':B':C'	830273.6	435081.7	277747.3

Table V. Energy Levels of H_2^{17}O (cm^{-1}).

1 ₀₁	23.773	5 ₀₅	324.658	7 ₀₇	584.941
1 ₁₁	36.931	5 ₁₅	325.878	7 ₁₇	585.162
1 ₁₀	42.187	5 ₁₄	398.877	7 ₁₆	702.888
2 ₀₂	70.004	5 ₂₄	415.126	7 ₂₆	708.022
2 ₁₂	79.226	5 ₂₃	445.792	7 ₂₅	781.381
2 ₁₁	94.970	5 ₃₃	502.179	7 ₃₅	814.616
2 ₂₁	134.144	5 ₃₂	507.174	7 ₃₄	840.868
2 ₂₀	135.430	5 ₄₂	607.165	7 ₄₄	924.647
3 ₀₃	136.536	5 ₄₁	607.403	7 ₄₃	928.301
3 ₁₃	141.901	5 ₅₁	737.629	7 ₅₃	1055.082
3 ₁₂	173.109	5 ₅₀	737.632	7 ₅₂	1055.283
3 ₂₂	205.479	6 ₀₆	445.717	8 ₀₈	742.401
3 ₂₁	211.434	6 ₁₆	446.243	8 ₁₈	742.494
3 ₃₁	283.559	6 ₁₅	541.996	8 ₁₇	881.108
3 ₃₀	283.765	6 ₂₅	551.610	8 ₂₇	883.664
4 ₀₄	221.618	6 ₂₄	601.961	8 ₂₆	981.504
4 ₁₄	224.302	6 ₃₄	647.074	8 ₃₆	1003.791
4 ₁₃	275.128	6 ₃₃	659.988	8 ₃₅	1048.662
4 ₂₃	299.436	6 ₄₃	753.712	8 ₄₅	1119.484
4 ₂₂	315.076	6 ₄₂	754.819	8 ₄₄	1128.935
4 ₃₂	380.804	6 ₅₂	884.104	8 ₅₄	1250.505
4 ₃₁	382.174	6 ₅₁	884.140	8 ₅₃	1251.298
4 ₄₁	485.209	6 ₆₁	1038.934		
4 ₄₀	485.237	6 ₆₀	1038.934		

in the literature. On the other hand, use of a theoretical model makes possible the introduction of information from other isotopic species and the analysis discussed above. The spectral constants which result from this analysis make possible the calculation of the set of energy levels shown in Table V.

One test of the accuracy of these energy levels is a comparison of them with combination difference energies calculated from the H_2^{17}O lines which were observed by Fraley, Rao, and Jones (6) and by Williamson and Rao (7). With the exception of a small number of combination differences, the agreement is excellent and comparable with the experimental uncertainty of the infrared data ($\sim 0.01 \text{ cm}^{-1}$). Most of these exceptions can be shown to be internally inconsistent with the other H_2^{17}O combination differences. In addition, there is no systematic deterioration in the agreement between the energy levels of Table V and the infrared combination differences over the range of J_r states covered by the combination differences. These results indicate that the uncertainties in the H_2^{17}O energy levels are similar to those derived for H_2^{16}O (17) and H_2^{18}O (6, 7). Since the accuracy of these levels depends upon the information contents transferred to the H_2^{17}O analysis from H_2^{16}O and H_2^{18}O analyses in the form of distortion constants, Table V includes only those energy levels for which the corresponding levels in H_2^{16}O and H_2^{18}O are well established.

ACKNOWLEDGMENT

The authors would like to thank Professor Walter Gordy for his encouragement and support during this work.

RECEIVED: November 20, 1974

REFERENCES

1. For an extensive bibliography see F. C. DE LUCIA, P. HELMINGER, AND W. H. KIRCHHOFF, *J. Phys. Chem. Ref. Data* **3**, 211 (1974).
2. G. STEENBECKELIERS AND J. BELLET, *Compt. Rend.* **273B**, 471 (1971).
3. P. HELMINGER, F. C. DE LUCIA, AND W. GORDY, *Phys. Rev. Lett.* **25**, 1397 (1970).
4. F. C. DE LUCIA, P. HELMINGER, R. L. COOK, AND W. GORDY, *Phys. Rev. A* **5**, 487 (1972).
5. F. C. DE LUCIA, P. HELMINGER, R. L. COOK, AND W. GORDY, *Phys. Rev. A* **6**, 1324 (1972).
6. P. E. FRALEY, K. NARAHARI RAO, AND L. H. JONES, *J. Mol. Spectrosc.* **29**, 312 (1969).
7. J. G. WILLIAMSON AND K. NARAHARI RAO, *J. Mol. Spectrosc.* **40**, 372 (1971).
8. F. C. DE LUCIA, *J. Mol. Spectrosc.*, to be published.
9. J. VERHOEVEN, A. DYMANUS, AND H. BLUYSSSEN, *J. Chem. Phys.* **50**, 3330 (1969).
10. F. C. DE LUCIA, P. HELMINGER, W. GORDY, H. W. MORGAN, AND P. A. STAATS, *Phys. Rev. A* **6**, 2785 (1973).
11. P. HELMINGER, F. C. DE LUCIA, W. GORDY, P. A. STAATS, AND H. W. MORGAN, *Phys. Rev. A* **10**, 1072 (1974).
12. J. K. G. WATSON, *J. Chem. Phys.* **45**, 1360 (1966).
13. J. K. G. WATSON, *J. Chem. Phys.* **46**, 1935 (1967).
14. J. K. G. WATSON, *J. Chem. Phys.* **48**, 4517 (1968).
15. D. KIVELSON AND E. B. WILSON, *J. Chem. Phys.* **20**, 1575 (1952).
16. W. GORDY AND R. L. COOK, "Microwave Molecular Spectra," Wiley, New York, 1970.
17. P. E. FRALEY AND K. NARAHARI RAO, *J. Mol. Spectrosc.* **29**, 348 (1969).