

Enthalpy of Formation of ${}^2\Pi_{3/2}$ SH

Attila G. Császár*

Department of Theoretical Chemistry, Eötvös University, H-1518 Budapest 112, P.O. Box 32, Hungary

Matthew L. Leininger

Sandia National Laboratory, MS 9217, Livermore, California

Alexander Burcat

Faculty of Aerospace Engineering, Technion—Israel Institute of Technology, Haifa 32000, Israel

Received: July 25, 2002; In Final Form: January 2, 2003

The standard enthalpy of formation, $\Delta_f H^\circ$, of ${}^2\Pi_{3/2}$ SH has been determined at converged levels of ab initio electronic structure theory, including high-order coupled cluster and full configuration interaction benchmarks. The atomic Gaussian basis sets employed include the (aug)-cc-p(wC)VnZ family with $n = 3, 4, 5$, and 6. Extrapolations to the complete one-particle basis set and the full configuration interaction limits, where appropriate, have been performed to reduce remaining computational errors. Additional improvements in the enthalpy of formation of ${}^2\Pi$ SH were achieved by appending the valence-only treatment with core–valence correlation, scalar relativistic and spin–orbit effects, and the diagonal Born–Oppenheimer correction. The recommended values for $\Delta_f H_0^\circ$ and $\Delta_f H_{298}^\circ$ of ${}^2\Pi$ SH are $141.24_{-0.46}^{+0.52}$ and $141.87_{-0.46}^{+0.52}$ kJ mol $^{-1}$, respectively, corresponding to a recommended $D_e = 365.76_{-0.42}^{+0.50}$ kJ mol $^{-1}$. The corresponding enthalpy of formation of ${}^2\Pi$ SD is $\Delta_f H_0^\circ = 140.17_{-0.50}^{+0.56}$ kJ mol $^{-1}$.

1. Introduction

H₂S is the major source of sulfur compounds in the earth's atmosphere. This compound is generated naturally by volcanic activity, by bacteria, and mainly by man-made pollution due to the burning of fossil fuels containing large amounts of S. The major source of the mercapto radical, SH, in the earth's atmosphere is S–H bond fission following near-ultraviolet photoexcitation of H₂S. Thermochemical data for SH are of special importance because of the role SH plays as an intermediate in reactions generating sulfur-containing pollutants (e.g., acid rain) as well as in complex reaction mechanisms explaining fossil combustion processes.¹

New, ever more efficient techniques for the solution of the electronic structure problem coupled with advances in computer architecture make possible, with unprecedented accuracy,^{2–11} the theoretical reevaluation of thermochemical properties. Because of their crucial importance in reaction kinetics studies, temperature-dependent standard enthalpies of formation, $\Delta_f H_7^\circ$, are of the greatest interest. For the smallest radicals, the accuracy of the best ab initio computations (see, for example, refs 10 and 11) of $\Delta_f H_7^\circ$ can exceed that of traditional experimental determinations.^{12–18}

The most straightforward way to determine the converged ab initio $\Delta_f H_0^\circ$ of ${}^2\Pi_{3/2}$ SH goes through the determination of its atomization energy, though there are more efficient computational routes for systems with more atoms. The best theoretical scheme for this study is offered by the focal-point approach.^{19,20} The converged ab initio determination of the enthalpy of formation of ${}^2\Pi_{3/2}$ SH is made simpler by the relatively small size of the system, allowing for the utilization of full config-

uration interaction (FCI) techniques.²¹ The calculations reported herein for the SH radical are the most extensive reported to date, though calculations with a similar purpose have been performed before.^{22–25}

Major thermochemical tables and databanks recommend values for the atomization energy, $D_{0/e}$, and for $\Delta_f H_7^\circ$ of ${}^2\Pi_{3/2}$ SH. The best available value of the zero-point-corrected atomization energy of ${}^2\Pi_{3/2}$ SH is $D_e = 350.51 \pm 1.20$ kJ mol $^{-1}$,²⁶ obtained from a careful experimental and theoretical analysis of the secondary photolysis of ground-state SH arising from the 218.2-nm photodissociation of H₂S. Note that this value corresponds to the $v'' = 0, J'' = 3/2$ level of the X ${}^2\Pi_{3/2}$ spin–orbit component and the lowest-energy products, S(³P₂) and H(²S). The available enthalpies of formation^{12,13,16,27–38} are summarized in Table 1. The available values for $\Delta_f H_{298}^\circ$ (SH) scatter around 140 kJ mol $^{-1}$, with recent reviews (e.g., ref 37) settling at 139.3 ± 3.0 kJ mol $^{-1}$. A slightly higher mean value, 143.0 ± 3.0 kJ mol $^{-1}$, has also been suggested and accepted¹³ on the basis of the high-quality work of Nicovich et al.³¹ Both values are in accord with the value obtained by Nourbakhsh and co-workers.³⁰ Nevertheless, in the same study, Nourbakhsh et al. suggested that $\Delta_f H_0^\circ = 148.5 \pm 8.4$ kJ mol $^{-1}$ for the enthalpy of formation of CH₃S, which is substantially different from the generally accepted low value of Nicovich,³¹ 131.5 ± 2.3 kJ mol $^{-1}$. To resolve this discrepancy, it is also mandatory to have a highly dependable estimate of the enthalpy of formation of SH.

2. Computational Approach

The X ${}^2\Pi$ ground electronic state of the mercapto radical has a $(1\sigma)^2(2\sigma)^2(3\sigma)^2(1\pi)^4(4\sigma)^2(5\sigma)^2(2\pi)^3$ electronic configuration. Single-reference methods are appropriate for the descrip-

* Corresponding author. E-mail: csaszar@chem.elte.hu.

TABLE 1: Literature Data for the Enthalpy of Formation of ${}^2\Pi$ SH^a

$\Delta_f H_{298}^{\circ}$	reference	year	method ^b	comments
Measurements				
144.76 ± 16.7	Mackle ²⁷	1963	KE	<i>c</i>
140.58 ± 4.6	Hwang and Benson ²⁸	1977	KE	<i>d</i>
138.6 ± 0.4	Traeger ²⁹	1984	PIMS-IE	<i>e</i>
142.4 ± 6.3	Nourbakhsh et al. ³⁰	1991	PIMS-IE	<i>f</i>
143.0 ± 2.8	Nicovich et al. ³¹	1992	KE	<i>g</i>
Computations				
141.0	Curtiss et al. ³²	1998	G3	<i>h</i>
138.0	Curtiss et al. ³³	1999	AI	<i>i</i>
140.6	Parthiban and Martin ³⁴	2001	W2	<i>j</i>
137.6	Janoschek and Rossi ³⁵	2002	G3MP2B3	<i>k</i>
143.1	Peebles and Marshall ²⁵	2002	AI	
Reviews and Evaluations				
149 ± 12	Kerr ³⁸	1966	CDE	<i>l</i>
140.6 ± 4.6	McMillen and Golden ³⁶	1982	CDE	<i>m</i>
140.4 ± 3.5	Gurvich et al. ¹²	1989	CDE	
143.0 ± 3.0	Berkowitz et al. ¹³	1994	CDE	<i>n</i>
139.3 ± 3.0	Chase et al. ¹⁶	1998	CDE	<i>o</i>
139.3 ± 5.0	Burcat ³⁷	2001	CDE	<i>p</i>

^a All values are in kJ mol⁻¹. ^b KE = kinetic equilibrium study. PIMS-IE = photoionization mass spectrometry- ionization energy determination. AI = ab initio computation using a model chemistry. CDE = critical data evaluation. ^c Average of three independent measurements: the dissociations of H₂S → HS + H, the ionization potential of HS, and the appearance potential of HS⁺ from H₂S. ^d Calculated from H₂S + I₂ reaction kinetics in the 555–595 K range. ^e PIMS has been used to measure the appearance energies of [C₂H₅]⁺ from ethanethiol, [C₃H₇]⁺ from 2-propanethiol, and [C₃H₅]⁺ from 2-methylthiirane. ^f $\Delta_f H_0^{\circ} = 141.8 \pm 6.3$ kJ mol⁻¹ was deduced from time-of-flight measurements of CH₃ and SH photofragments from the photodissociation of CH₃SH in a supersonic molecular beam. ^g Kinetics of the reaction Br(²P_{3/2}) + H₂S → SH + HBr using resonance fluorescence temperature-dependent experiments result in $\Delta_f H_0^{\circ} = 142.55 \pm 3.01$ kJ mol⁻¹ and $\Delta_f H_{298}^{\circ} = 143.01 \pm 2.85$ kJ mol⁻¹. ^h G2 and G3 model chemistry values are both reported. ⁱ G3(MP2/6-31G(d)) computation. ^j Weizmann model W2. The W1 value is 145.6 kJ mol⁻¹. ^k The estimated average error in the G3MP2B3 calculations is 8 kJ mol⁻¹. ^l The old value of Kerr is an uncertain estimate based on the dissociation energy of the H–S bond as 90 ± 2 kcal/mol from the pyrolysis of benzyl sulfide. ^m Refers to the value of Hwang and Benson.²⁸ ⁿ Extensive calculation of data (mainly kinetic determinations) published until 1992. The recommended value is that of Nicovich et al.³¹ ^o Based on a photoionization appearance energy (AE) of SH⁺ produced from H₂S (Dibeler, V. H.; Liston, S. K. *J. Chem. Phys.* **1968**, *49*, 482) and the spectroscopic ionization energy for the SH radical (Morrow, B. A. *Can. J. Phys.* **1966**, *44*, 2447). These data were not reviewed after 1977. As pointed out by Traeger,²⁹ using a better estimate for the AE of SH⁺ (14.300 ± 0.0024 eV; Prest, H. F.; Tzeng, W.-B.; Brom, J. M., Jr.; Ng, C. Y. *Int. J. Mass Spectrom. Ion Phys.* **1983**, *50*, 315) increases $\Delta_f H_{298}^{\circ}$ to 142 ± 3 kJ mol⁻¹, in perfect agreement with our recommended (computational) value of 141.87^{+0.52}_{-0.46} kJ mol⁻¹. ^p Thermochemical database for combustion. The old calculation was based on 1977 JANAF¹⁶ tables.

tion of the electronic structures of H, S, and SH. Because electronic energy calculations based on the correlation-consistent (cc) family of Gaussian basis sets (aug)-cc-p(C)VXZ (with cardinal number $X = 2(D), 3(T), 4(Q), 5$, and 6) of Dunning and co-workers^{39–42} usually approach the complete basis set limit in a systematic fashion, these basis sets were employed in the focal-point basis set extrapolations of the present study. Nevertheless, several problems with the original correlation-consistent basis sets for S have been observed (see, for example, refs 43 and 44). Convergence problems at the Hartree–Fock level led to the development of a new standard set of cc basis sets, denoted (aug)-cc-pV(X+d)Z,^{41,42} which have been employed in this study instead of the original cc basis sets. Since no core-correlated basis sets are available in ref 41, CVXZ basis sets, with $X = 4, 5$, and 6, were constructed⁴⁵ as an even-tempered extension, with a factor of 3, of the corresponding completely uncontracted cc-pVXZ sets by adding two tight functions to each shell beyond p. A lack of optimization of the exponents in the CVXZ basis sets leads to convergence problems. Therefore, an unpublished set⁴⁶ of polarization-weighted basis sets for S has also been employed, denoted cc-pwCVXZ, with $X = 3, 4$, and 5. Complete basis set limit energies have been estimated through the equations $E^X = E_{\text{limit}} + a \exp(-bX)$ ⁴⁷ and $E^X = E_{\text{limit}} + cX^{-3}$ ⁴⁸ for the HF and correlation energy limits, respectively, employing the best three and two energies available, in order. Reference electronic wave functions have been determined by the single-configuration restricted (open-shell) Hartree–Fock [R(O)HF] method. Dynamical electron correlation was accounted for by the coupled cluster (CC) method including all single and double (CCSD)⁴⁹

and triple excitations (CCSDT).⁵⁰ The CCSD(T) method,^{51,52} which estimates the effect of connected triple excitations through a perturbative term [(T)], was also employed extensively. The full configuration interaction (FCI) computations²¹ utilized an ROHF reference wave function. During valence correlation energy computations, the 1s, 2s, and 2p core orbitals of sulfur were excluded from the active space. No virtual orbitals were frozen in any of the calculations. The experimental equilibrium bond distance $r_e/\text{\AA} = 1.3409$ ⁵³ was adopted for all electronic structure computations in the valence focal-point analysis and during the auxiliary energy calculations. Core-correlation effects were determined by means of all-electron and frozen-core treatments up to CCSDT with (pw)CVXZ ($X = 3, 4$, and 5) and CVXZ ($X = 4, 5$, and 6) basis sets. Scalar relativistic effects^{54,55} were gauged by first-order perturbation theory applied to the one-electron mass–velocity and Darwin terms (MVD1). The computation of the diagonal Born–Oppenheimer (DBOC) correction^{56,57} was performed at the Hartree–Fock level within the formalism of Handy, Yamaguchi, and Schaefer.⁵⁶ Different versions of the program packages ACES II⁵⁸ and PSI⁵⁹ were used for the electronic structure computations.

The total energies computed as part of this study for H(²S), S(³P), and SH (²Π) are given in Tables 4S–6S of the Supporting Information.

3. Discussion

Tables 2 and 3 summarize the atomization energy results obtained using the focal-point scheme^{19,20} for the valence-only and core–valence treatments, respectively. The symbol δ in

TABLE 2: Valence Focal-Point Analysis of the Dissociation Energy of ${}^2\Pi$ SH in kJ mol^{-1a}

basis	ROHF	$\delta[\text{CCSD}]$	$\delta[\text{CCSD(T)}]$	$\delta[\text{CCSDT}]$	$\delta[\text{FCI}]$	FCI
aug-cc-pV(D+d)Z(9/32)	252.12	+84.47	+3.67		+0.18	340.77
cc-pwCVTZ(14/59)	254.76	+95.23	+3.87			
aug-cc-pV(T+d)Z(23/55)	255.82	+97.65	+4.17	+0.09	+0.18 ^b	357.91
cc-pwCVQZ(30/109)	256.32	+101.53	+4.23	+0.01		
aug-cc-pV(Q+d)Z(46/89)	256.39	+102.44	+4.30	+0.07	+0.18 ^b	363.38
CVQZ(30/134)	256.34	+101.76	+4.25	+0.01		
cc-pwCV5Z(55/181)	256.57	+103.30	+4.37			
aug-cc-pV(5+d)Z(80/136)	256.59	+103.99	+4.47			
CV5Z(55/185)	256.54	+103.31	+4.37			
aug-cc-pV(6+d)Z(127/185)	256.68	+104.19	+4.48			
CV6Z(91/273)	256.65	+104.03	+4.44			
extrapolated[4–6]	256.73	+104.47	+4.49	+0.05	+0.18	365.92

^a See text for an explanation of column headings and basis sets and for details about the extrapolation to the basis set limit for ROHF, $\delta[\text{CCSD}]$, and $\delta[\text{CCSD(T)}]$. After each basis set, the number of contracted Gaussian basis functions for H/S is given in parentheses. Reference geometry: $r(\text{S}-\text{H}) = 1.3409 \text{ \AA}$. ^b Obtained using SHOC factors (see text for details).

TABLE 3: Core–Valence Correlation Corrections (kJ mol^{-1}) to the Total Atomization Energy of ${}^2\Pi$ SH^a

basis	$\delta[\text{CCSD}]$	$\delta[\text{CCSD(T)}]$	$\delta[\text{CCSDT}]$	$\Delta E_c(\text{CC})$
cc-pwCVTZ	+0.615	+0.291	+0.013	+0.919
cc-pwCVQZ	+0.493	+0.304	+0.021	+0.818
cc-pwCV5Z	+0.472	+0.307		
extrapolated[4,5]	+0.450	+0.310	[+0.021]	+0.781
CVQZ	+0.447	+0.298	+0.022	+0.767
CV5Z	+0.559	+0.301		

^a See text for an explanation of column headings and basis sets and for details about the extrapolation to the basis set limit for ROHF, $\delta[\text{CCSD}]$, and $\delta[\text{CCSD(T)}]$. Reference geometry: $r(\text{S}-\text{H}) = 1.3409 \text{ \AA}$.

these Tables denotes the increment in the relative energy (ΔE_c) with respect to the preceding level of theory, as given by the hierarchy ROHF \rightarrow CCSD \rightarrow CCSD(T) \rightarrow CCSDT \rightarrow FCI.

In a recent study,⁶⁰ a simple multiplicative procedure, termed scaled higher-order correlation or SHOC, was suggested to estimate higher-order correlation (HOC) energies not covered, for example, in CCSD(T) or CCSDT treatments. This procedure utilizes the observation that HOC energy increments show limited basis set dependence and thus even at the complete basis set limit they can be estimated from explicit small basis set FCI and CCSD(T) or CCSDT calculations. From our extensive previous computations, it became clear that the augmented basis sets are more amenable for this correction. The aug-cc-pVXZ CCSDT SHOC scale factors for $\text{S}({}^3\text{P})$ are 1.001508 and 1.001124 for $X = 2$ and 3, respectively. The aug-cc-pVDZ CCSDT SHOC scale factor for $\text{SH}({}^2\Pi)$ is 1.001634. As expected, it is very similar to the corresponding $\text{S}({}^3\text{P})$ SHOC factor. The SHOC factors at the CCSD(T) level are substantially larger; using the aug-cc-pV(D+d)Z basis set, they are 1.005967 and 1.005926 for $\text{S}({}^3\text{P})$ and $\text{SH}({}^2\Pi)$, respectively. To make a balanced treatment, the CCSDT SHOC factors of 1.001124 and 1.001250 were employed for the valence-only treatments of $\text{S}({}^3\text{P})$ and $\text{SH}({}^2\Pi)$, respectively, where the latter factor was obtained by scaling the former one by 1.001634/1.001508. Using the best CCSD(T) and CCSDT sets of SHOC factors shows agreement within 0.1 kJ mol^{-1} for the atomization energy of ${}^2\Pi$ SH.

The directly computed valence-only atomization energies provide lower limits to the correct result, as an extension of the basis set as well as the electron correlation treatment increases, as expected, the calculated atomization energies. The valence-only complete basis set ROHF atomization energy is 256.73 \pm 0.04 kJ mol^{-1} , obtained after averaging the extrapolated aug-cc-pVXZ and CVXZ, $X = 4, 5$, and 6 estimates and multiplying their deviation from the average by 2 to attach an

error estimate to the recommended value. Estimating $\delta[\text{CCSD}]$ at the complete basis set limit presents, as usual, some difficulty. Nevertheless, the complete basis set CCSD value obtained from the aug-cc-pVXZ, $X = 5$ and 6, basis set results differs from the aug-cc-pV(6+d)Z and CV6Z values by only 0.28 and 0.44 kJ mol^{-1} , respectively, which are comfortably small. Our best estimate for $\delta[\text{CCSD}]$ is $+104.47_{-0.15}^{+0.30} \text{ kJ mol}^{-1}$. The extrapolated CCSD(T) increment, obtained the same way, is $+4.49_{-0.01}^{+0.05} \text{ kJ mol}^{-1}$. It is impossible to estimate the $\delta[\text{CCSDT}]$ and $\delta[\text{FCI}]$ increments accurately from the limited data available. Our best estimate for their combined effect is $+0.23 \pm 0.10 \text{ kJ mol}^{-1}$. Therefore, the valence-only complete basis set FCI atomization energy of ${}^2\Pi_{3/2}$ SH (see Table 2) is $365.92_{-0.19}^{+0.32} \text{ kJ mol}^{-1}$. As seen in Table 3, the core contribution to the atomization energy of $\text{SH}({}^2\Pi)$ is rather small. The best estimate is $+0.78 \pm 0.15 \text{ kJ mol}^{-1}$ at the estimated complete basis set FCI limit, obtained from the extrapolated cc-pwCVXZ, $X = 4$ and 5 results. It is notable how different the convergence characteristics of the CVXZ and cc-pwCVXZ $\delta[\text{CCSD}]$ results are. This is most likely due to weaknesses in our design of the CVXZ basis sets.

The scalar relativistic correction to the atomization energy of ${}^2\Pi$ SH has been computed at the HF and CCSD(T) levels using the pwCVQZ basis set and the MVD1 formalism. The correction is $-0.85 \text{ kJ mol}^{-1}$. The correlation contribution included in this value is $+0.23 \text{ kJ mol}^{-1}$. The fact that two-electron relativistic energy corrections are usually comparable to the electron correlation contribution to the MVD1 energy correction⁵⁵ indicates that relativistic corrections beyond MVD1 should have a small effect even at the level of precision sought in this study. Therefore, we set the scalar relativistic correction to $-0.85 \pm 0.30 \text{ kJ mol}^{-1}$. SH and S have inverted ground electronic states, ${}^2\Pi_{3/2}$ and ${}^3\text{P}_2$, respectively. For open-shell states, traditional nonrelativistic electronic structure calculations yield the weighted average of the available multiplets. To obtain the energy of the lowest states of S and SH, the next relativistic energy correction, the spin–orbit effect must be considered. We are employing literature values for these quantities; the experimental weighted-average spin–orbit splitting constants are 2.345 kJ mol^{-1} for the ${}^3\text{P}$ state of sulfur^{61,62} and 2.254 kJ mol^{-1} for SH.⁶³ The spin–orbit correction to the dissociation energy is thus $-0.091 \pm 0.010 \text{ kJ mol}^{-1}$. Therefore, the total relativistic correction is $-0.94 \pm 0.30 \text{ kJ mol}^{-1}$, to be added to the computed nonrelativistic atomization energy.

The DBOC corrections, in cm^{-1} , computed at the Hartree–Fock level using a medium-sized TZ2P+diff basis set are +59.65, +1243.03, and +1302.90 for $\text{H}({}^2\text{S})$, $\text{S}({}^3\text{P})$, and $\text{SH}({}^2\Pi)$, respectively, resulting in an overall DBOC correction to the

atomization energy of 0.2 cm^{-1} . Therefore, this correction is negligible for this study. Collecting all of the terms and assuming independent error bars for the different terms, we obtain our best estimate for the equilibrium atomization energy of ${}^2\Pi_{3/2}$ SH as $365.92_{-0.19}^{+0.32} + 0.78_{-0.15}^{+0.15} - 0.94_{-0.30}^{+0.30} = 365.76_{-0.39}^{+0.46} \text{ kJ mol}^{-1}$. The molecular zero-point energy (ZPE) of ${}^{32}\text{S}^1\text{H}$ is $16.04 \text{ kJ mol}^{-1}$,⁵³ obtained as $(1/2)\omega_e - (1/4)\omega_e x_e$ with $\omega_e = 2711.6 \text{ cm}^{-1}$ ⁵³ and $\omega_e x_e = 60 \text{ cm}^{-1}$.⁶⁴ The error bar that can be attached to this quantity is perhaps $\pm 0.20 \text{ kJ mol}^{-1}$. This brings the computed zero-point-corrected atomization energy, corresponding to the lowest-energy spin multiplet, to $349.72_{-0.44}^{+0.50} \text{ kJ mol}^{-1}$. This value is in agreement with (a) the best experimental values of $350.51 \pm 1.20 \text{ kJ mol}^{-1}$ ²⁶ and $349.4 \pm 2.9 \text{ kJ mol}^{-1}$,⁶⁵ though it has a considerably lower error bar and (b) the best previous direct theoretical estimate of $D_0(\text{S}-\text{H}) = 349.9 \pm 2.0 \text{ kJ mol}^{-1}$ ²⁵ based principally on coupled-cluster computations. (Half of the difference between the computed mean D_0 values is due to the use of slightly different ZPE corrections.)

The calculation of the enthalpy of formation from the atomization energy requires a knowledge of the enthalpy of formation of atoms H and S in their respective ground states. The relevant data, when available, were taken from ref 14 and are given, in kJ mol^{-1} , as $\Delta_f H_{298}^\circ[\text{S}(^3\text{P})] = 277.17 \pm 0.15$, $\Delta_f H_{298}^\circ[\text{H}(^2\text{S})] = 217.998 \pm 0.006$, $H_{298} - H_0[\text{H}] = 6.197 \pm 0.001$, $H_{298} - H_0[\text{H}_2] = 8.468 \pm 0.001$, $H_{298} - H_0[\text{S}_{\text{cr, rhombic}}] = 4.412 \pm 0.006$, and $H_{298} - H_0[\text{S}_{\text{gas}}] = 6.657 \pm 0.001$. The resulting 0 K atomic enthalpies of formation are $\Delta_f H_0^\circ[\text{H}(^2\text{S})] = 216.034 \pm 0.006 \text{ kJ mol}^{-1}$ and $\Delta_f H_0^\circ[\text{S}(^3\text{P})] = 274.925 \pm 0.150 \text{ kJ mol}^{-1}$. The $H_{298} - H_0$ value for SH was taken from ref 16 as $9.274 \text{ kJ mol}^{-1}$. On the basis of these experimental values, the final computational value for $\Delta_f H_0^\circ$ of ${}^2\Pi_{3/2}$ SH is $141.24_{-0.46}^{+0.52} \text{ kJ mol}^{-1}$. The best recommended computational value for $\Delta_f H_{298}^\circ$ is $141.87_{-0.46}^{+0.52} \text{ kJ mol}^{-1}$. Our mean value is significantly higher than $139.3 \pm 3.0 \text{ kJ mol}^{-1}$ recommended by JANAF¹⁶ and somewhat lower than $143.0 \pm 3.0 \text{ kJ mol}^{-1}$ recommended in other databases¹³ on the basis of the work of Nicovich et al.³¹ Our final recommended values are also significantly different from the $\Delta_f H_0^\circ(\text{SH})$ and $\Delta_f H_{298}^\circ(\text{SH})$ values recommended in ref 25, $142.9 \pm 0.8 \text{ kJ mol}^{-1}$ and $143.1 \pm 0.8 \text{ kJ mol}^{-1}$, respectively, on the basis of a certain combination of computed and measured energy values.

4. Conclusions

In this paper, we have calculated ab initio the equilibrium and zero-point-energy-corrected dissociation energies of ${}^2\Pi_{3/2}$ SH with recommended values of $365.76_{-0.39}^{+0.46}$ and $349.72_{-0.44}^{+0.50} \text{ kJ mol}^{-1}$, respectively. These results allow us to estimate the standard enthalpy of formation, $\Delta_f H_T^\circ$ of ${}^2\Pi_{3/2}$ SH. The results obtained, $141.24_{-0.46}^{+0.52}$ and $141.87_{-0.46}^{+0.52} \text{ kJ mol}^{-1}$ for $\Delta_f H_0^\circ$ and $\Delta_f H_{298}^\circ$, respectively, are somewhat different from the best available experimental results and have smaller error bars. Our final results for the enthalpy of formation of ${}^2\Pi_{3/2}$ SH shows that we can reliably calculate $\Delta_f H_T^\circ$ for such small species within about 0.5 kJ mol^{-1} .

To get a converged theoretical estimate of the enthalpy of formation of ${}^2\Pi$ SD, only a few extra data points are necessary over the benchmark-quality results obtained for SH. The difference in the zero-point energies of SH and SD is $4.85 \pm 0.20 \text{ kJ mol}^{-1}$ based on $\omega_e = 1885.6 \text{ cm}^{-1}$ ⁵³ and $\omega_e x_e = 31 \text{ cm}^{-1}$ ⁶⁴ for SD. The difference in the spin-orbit splitting constants of SH and SD¹⁸ is only $0.001 \text{ kJ mol}^{-1}$, which is negligible for this study. Since the DBOC correction was found

to be negligible for SH, the same can safely be assumed for SD. The enthalpy of formation of D(${}^2\text{S}$) can be obtained from standard tables: $\Delta_f H_0^\circ[\text{D}(^2\text{S})] = 219.81 \text{ kJ mol}^{-1}$. Therefore, we obtain the result $\Delta_f H_0^\circ[\text{SD}(^2\Pi)] = 140.17_{-0.50}^{+0.56} \text{ kJ mol}^{-1}$.

Acknowledgment. The work of A.G.C. has been supported by the Scientific Research Fund of Hungary (OTKA T033074). M.L.L. was supported by Sandia National Laboratories. Sandia is a multiprogram laboratory operated by Sandia Corporation, a Lockheed Martin Company, for the United States Department of Energy under contract DE-AC04-94AL85000. The research described forms part of an effort by a task group of the International Union of Pure and Applied Chemistry (2000-013-2-100) to determine structures, vibrational frequencies, and thermodynamic functions of free radicals of importance in atmospheric chemistry.

Supporting Information Available: ROHF energies for H(${}^2\text{S}$), valence-only and all-electron energies for S(${}^3\text{P}$), and valence-only and all-electron energies for ${}^2\Pi$ SH. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes

- Hynes, A. J.; Wine, P. H. *Kinetics and Mechanism of the Oxidation of Gaseous Sulfur Compounds*. In *Gas-Phase Combustion Chemistry*; Gardiner, W. C., Jr., Ed.; Springer-Verlag, New York, 2000; Chapter 3, p 343.
- East, A. L. L.; Allen, W. D. *J. Chem. Phys.* **1993**, *99*, 4638.
- Ruscic, B.; Feller, D.; Dixon, D. A.; Peterson, K. A.; Harding, L. B.; Asher, R. L.; Wagner, A. F. *J. Phys. Chem. A* **2001**, *105*, 1.
- Dixon, D. A.; Feller, D.; Peterson, K. A. *J. Chem. Phys.* **2001**, *115*, 2576.
- Dixon, D. A.; Feller, D.; Sandrone, G. *J. Phys. Chem. A* **1999**, *103*, 4744.
- Martin, J. M. L.; de Oliveira, G. *J. Chem. Phys.* **1999**, *111*, 1843.
- Irikura, K. K.; Frurip, D. J. *Computational Thermochemistry: Prediction and Estimation of Molecular Thermodynamics*; American Chemical Society: Washington, DC, 1998.
- Feller, D.; Dixon, D. A. *J. Phys. Chem. A* **2000**, *104*, 3048.
- Feller, D. *J. Comput. Chem.* **1996**, *17*, 1571; the database is available at <http://www.emsl.pnl.gov:2080/proj/crdb/>.
- Ruscic, B.; Wagner, A. F.; Harding, L. B.; Asher, R. L.; Feller, D.; Dixon, D. A.; Peterson, K. A.; Song, Y.; Qian, X.; Ng, C.-Y.; Liu, J.; Chen, W.; Schwenke, D. W. *J. Phys. Chem. A* **2002**, *106*, 2727.
- Császár, A. G.; Szalay, P. G.; Leininger, M. L. *Mol. Phys.* **2002**, *100*, 3879.
- Gurvich, L. V.; Veyts, I. V.; Alcock, C. B. *Thermodynamic Properties of Individual Substances*, 4th ed.; Hemisphere: New York, 1989; Vols. 1 and 2.
- Berkowitz, J.; Ellison, G. B.; Gutman, D. *J. Phys. Chem.* **1994**, *98*, 2744.
- Cox, J. D.; Wagman, D. D.; Medvedev, V. A. *CODATA Key Values for Thermodynamics*; Hemisphere: New York, 1989; see also <http://www.codata.org/codata/databases/key1.html>.
- DeMore, W. B.; Sander, S. P.; Golden, D. M.; Hampson, R. F.; Kurylo, M. J.; Howard, C. J.; Ravishankara, A. R.; Kolb, C. E.; Molina, M. J. *Chemical Kinetics and Photochemical Data for Use in Stratospheric Modeling*; Evaluation No. 12, JPL Publication 97-4, NASA and JPL, 1997.
- (a) Chase, M. W., Jr. *NIST-JANAF Thermochemical Tables*, 4th ed.; *J. Phys. Chem. Ref. Data* **1998**; Monograph No. 9; NIST WebBook: <http://webbook.nist.gov/>. (b) Chase, M. W., Jr.; Curnutt, J. L.; Downey, J. R.; McDonald, R. A.; Syverud, A. N.; Valenzuela, E. A. *J. Phys. Chem. Ref. Data* **1982**, *11*, 695.
- Kerr, J. A.; Stocker, D. W. In *Handbook of Chemistry and Physics*, 81st ed.; Lide D. R., Ed.; CRC Press: Boca Raton, FL, 2000–2001.
- Computational Chemistry Comparison and Benchmark Database, <http://srdata.nist.gov/cccbdb/>.
- East, A. L. L.; Allen, W. D.; Császár, A. G. In *Structures and Conformations of Non-Rigid Molecules*; Laane, J.; Dakkouri, M.; van der Veken, B.; Oberhammer, H., Eds.; Kluwer: Dordrecht, The Netherlands, 1993.
- Császár, A. G.; Allen, W. D.; Schaefer, H. F., III. *J. Chem. Phys.* **1998**, *108*, 9751.
- Sherrill, C. D.; Schaefer, H. F., III. The Configuration Interaction Method: Advances in Highly Correlated Approaches. In *Advances in*

Quantum Chemistry; Löwdin, P.-O., Ed.; Academic Press: New York, 1999; Vol. 34, p 143.

- (22) Hirst, D. M.; Guest, M. F. *Mol. Phys.* **1982**, *46*, 427.
(23) Bruna, P. J.; Hirsch, G. *Mol. Phys.* **1987**, *61*, 1359.
(24) Park, J. K.; Sun, H. S. *Chem. Phys. Lett.* **1992**, *195*, 469.
(25) Peebles, L. R.; Marshall, P. J. *J. Chem. Phys.* **2002**, *117*, 3132.
(26) Morley, J. *Chem. Soc., Faraday Trans.* **1993**, *89*, 3865.
(27) Mackle, H. *Tetrahedron* **1963**, *19*, 1159.
(28) Hwang, R. J.; Benson, S. W. *Int. J. Chem. Kinet.* **1977**, *11*, 579.
(29) Traeger, J. C. *Org. Mass Spectrom.* **1984**, *19*, 514.
(30) Nourbakhsh, S.; Norwood, K.; Yin, H.-M.; Liao, C.-L.; Ng, C. Y. *J. Chem. Phys.* **1991**, *95*, 946.
(31) Nicovich, J. M.; Kreutter, K. D.; van Dijk, C. A.; Wine, P. H. *J. Phys. Chem.* **1992**, *96*, 2518.
(32) Curtiss, L. A.; Raghavachari, K.; Redfern, P. C.; Rassolov, V.; Pople, J. A. *J. Chem. Phys.* **1998**, *109*, 7764.
(33) Curtiss, L. A.; Redfern, P. C.; Raghavachari, K.; Rassolov, V.; Pople, J. A. *J. Chem. Phys.* **1999**, *110*, 4703.
(34) Parthiban, S.; Martin, J. M. L. *J. Chem. Phys.* **2001**, *114*, 6014.
(35) Janoschek, R.; Rossi, M. J. *Int. J. Chem. Kinet.* **2002**, *34*, 550.
(36) McMillan, D. F.; Golden, D. M. *Annu. Rev. Phys. Chem.* **1982**, *33*, 493.
(37) Burcat, A. *Third Millennium Ideal Gas and Condensed Phase Thermodynamical Database for Combustion*; TAE Report No. 867; Technion: Haifa, Israel, 2001.
(38) Kerr, J. A. *Chem. Rev.* **1966**, *66*, 465.
(39) Dunning, T. H., Jr. *J. Chem. Phys.* **1989**, *90*, 1007.
(40) Wilson, A. K.; van Mourik, T.; Dunning, T. H., Jr. *J. Mol. Struct.: THEOCHEM* **1997**, *388*, 339.
(41) The (aug)-cc-p(C)VXZ basis sets were obtained from the Extensible Computational Chemistry Environment Basis Set Database, version 1.0, as developed and distributed by the Molecular Science Computing Facility, Environmental and Molecular Sciences Laboratory, which is part of the Pacific Northwest Laboratory, P.O. Box 999, Richland, WA 99352, and is funded by the U.S. Department of Energy. The Pacific Northwest Laboratory is a multiprogram laboratory operated by Battelle Memorial Institute for the U.S. Department of Energy under contract DE-AC06-76RLO 1830.
(42) Dunning, T. H., Jr.; Peterson, K. A.; Wilson, A. K. *J. Chem. Phys.* **2001**, *114*, 9244.
(43) Tarczay, G.; Császár, A. G.; Leininger, M. L.; Klopper, W. *Chem. Phys. Lett.* **2000**, *322*, 119.
(44) Martin, J. M. L. *J. Chem. Phys.* **1998**, *108*, 2791, and references therein.
(45) Császár, A. G.; Allen, W. D. *J. Chem. Phys.* **1996**, *104*, 2746.
(46) Peterson, K. A. Personal communication, 2002.
(47) Feller, D. *J. Chem. Phys.* **1992**, *96*, 6104.
(48) Klopper, W.; Bak, K. L.; Jørgensen, P.; Olsen, J.; Helgaker, T. *J. Phys. B: At. Mol. Opt. Phys.* **1999**, *32*, 103.
(49) Purvis, G. D.; Bartlett, R. J. *J. Chem. Phys.* **1982**, *76*, 1910.
(50) Noga, J.; Bartlett, R. J. *J. Chem. Phys.* **1987**, *86*, 7041.
(51) Raghavachari, K.; Trucks, G. W.; Pople, J. A.; Head-Gordon, M. *Chem. Phys. Lett.* **1989**, *157*, 479.
(52) Bartlett, R. J.; Watts, J. D.; Kucharski, S. A.; Noga, J. *Chem. Phys. Lett.* **1990**, *165*, 513.
(53) Huber, K. P.; Herzberg, G. *Molecular Spectra and Molecular Structure*; Van Nostrand Reinhold Co.: New York, 1979; Vol. 4.
(54) Balasubramanian, K. *Relativistic Effects in Chemistry, Part A: Theory and Techniques and Part B: Applications*; Wiley & Sons: New York, 1997.
(55) Tarczay, G.; Császár, A. G.; Klopper, W.; Quiney, H. M. *Mol. Phys.* **2001**, *99*, 1769, and references therein.
(56) Handy, N. C.; Yamaguchi, Y.; Schaefer, H. F., III. *J. Chem. Phys.* **1986**, *84*, 4481.
(57) Kutzelnigg, W. *Mol. Phys.* **1997**, *90*, 909.
(58) Stanton, J. F.; Gauss, J.; Watts, J. D.; Nooijen, M.; Oliphant, N.; Perera, S. A.; Szalay, P. G.; Lauderdale, W. J.; Gwaltney, S. R.; Beck, S.; Balkova, A.; Bernhold, D. E.; Baeck, K.-K.; Rozyczko, P.; Huber, C.; Bartlett, R. J. *Aces II*. The package also contains modified versions of the MOLECULE Gaussian integral program of Almlöf, J. and Taylor, P. R., the ABACUS integral derivative program written by Helgaker, T. U.; Jensen, H. J. Aa.; Jørgensen, P.; and Taylor, P. R., and the PROPS property evaluation integral code of Taylor, P. R.
(59) Crawford, T. D.; Sherrill, C. D.; Valeev, E. F.; Fermann, J. T.; Leininger, M. L.; King, R. A.; Brown, S. T.; Janssen, C. L.; Seidl, E. T.; Yamaguchi, Y.; Allen, W. D.; Xie, Y.; Vacek, G.; Hamilton, T. P.; Kellogg, C. B.; Remington, R. B.; Schaefer, H. F., III. *Psi 3.0*; PSITECH Inc.: Watkinsville, GA, 1999.
(60) Császár, A. G.; Leininger, M. L. *J. Chem. Phys.* **2001**, *114*, 5491.
(61) Moore, C. E. *Atomic Energy Levels*; U.S. National Bureau of Standards Circular 467; National Bureau of Standards: Washington, DC, 1949.
(62) Bashkin, S.; Stoner, J. R. *Atomic Energy Levels and Grottrian Diagrams*, 2nd ed.; North-Holland: Amsterdam, 1978.
(63) Ashworth, S. H.; Brown, J. M. *J. Mol. Spectrosc.* **1992**, *153*, 31.
(64) Ramsey, D. A. *J. Chem. Phys.* **1952**, *20*, 1920.
(65) Continetti, R. E.; Balko, B. A.; Lee, Y. T. *Chem. Phys. Lett.* **1991**, *182*, 400, 88, 841.